
- "Everything rests by changing" Heraclitus, (535-475 BCE)





Single-Trial Variability
in Early Visual Neuromagnetic Responses:
An Explorative Study based on the Regional Activation
Contributing to the N70m peak

[NeuroImage. vol.20(2): 765-783, 2003]




- is N70m peak a reproducible response at the single-trial level?
- is "signal + noise" a valid model?
- is ensemble averaging an appropriate response-recovery technique?

Objectives


- to exploit the temporal resolution of MEG signal to precisely characterize the nature of response variability
 - to identify the dominant sources contributing to the variability observed in the MEG signal
 - to associate the variability seen in the multichannel signal with the activity in early visual areas


[F. Moradi et al., NeuroImage, 2003]

"Consistent and precise localization of brain activity in human primary visual cortex by MEG and fMRI"




From the multichannel MEG signal to tomographic estimates of brain activity















80 msec

MEG signal recording and preprocessing


- 2 subjects from the "fMRI & MEG"-study participated
- 3 different runs of pattern reversal stimulation (1.43 Hz) & 1 subject baseline run were recorded



- for each run, 240 trials were recorded, with 625 Hz s.f. using the 151-channel whole-head Omega biomagnetometer.
- 3rd gradient formation, band-pass filter: 1-120 Hz, powerline removal, cardiac artifact elimination.
- trials were extracted: -100 to 200 msec and those without eye related activity were selected.

Methods

① Design of the spatial filter used to extract the temporal patterns conveying the regional response dynamics



2 Pattern Analysis of the extracted ST-patterns

Feature extraction


Embedding in Feature Space Clustering & Vector Quantization


Minimal Spanning Tree of the codebook


Interactive Study of pattern variability


MST-ordering of the code vectors


Orderly presentation of response variability














3 within-group analysis of regional response dynamics





• within-group analysis of multichannel single-trial signals





within-group analysis of single-trial MFT-solutions






Results

The Virtual Sensor design from the averaged MEG signal (4.5°-run)


Feature extraction based on the N70m latency-range




the brain waves captured by the Virtual Sensor mainly reflect the dynamic behavior of the generators responsible for the formation of the averaged N70m-peak


Based on the single-trial snapshots of regional brain activity corresponding to the N70m-related latency-range the response variability was described using a Vector-Quantization scheme.

- → The design of the encoder
- The applique idesign of the encoder (from the 4.5° run, k=10)






The application of the encoder (to all runs)




Visualizing the VQ results


i. By computing a 2D representation of the designed codebook and ii. depicting graphically the distribution of N70m snapshots in the k=10 groups with the node size



The codebook could be kept fixed across the different runs




The Voronoi-regions
were populated
in a stimulus-size-dependent manner




By within-group averaging, **prototypes of regional brain activity** were formed.

The computed prototypes of evoked response were compared with each other and contrasted with the ensemble average



a rich dynamic behavior of the N70m generators was revealed

The 2D-sketch of the codebook




transient response vs. oscillatory activity


Distinct response modes?


The ordered prototypes of regional brain activity

Using two signal descriptors:

the *time-varying energy & the inter-trial synchronization* expressing *amplitude-modulation* and *phase-modulation* accordingly, we provided a within-group characterization of the evoked response







A highly adaptive system is stimulated

Within a run the response mode changes:


from

stimulus-locked *increase in the energy* (10) to


phase-resetting of the ongoing regional activity (5)


..... And stimulus-induced desynchronization

... or *Enhancement* of the ongoing rhythmic activity




Trial-to-trial non-stationarity











The transition from a high-responsiveness state to a low-responsiveness state was the most important "law" governing the sequence of responses






which are the generators of N70m-response variability?

We applied the *k-partition* to the corresponding single-trial estimates of current source density produced by tomographic analysis (*MFT*)


and


we computed spatiotemporal prototypes of localized brain activity

(MFT-based prototypes)




The k-partition was the "thread" connecting the MEG-signal with the tomographic estimates










65 msec




80 msec


By visual exploration of MFT-based prototypes:

- Polymodal parietal areas
(BA5 & BA7)
contribute mostly
to the N70m-related
brain waves


- V1 contribution is considerably weaker




Is there any relationship between N70m-response variability and activity in early visual areas?



- structural similarity
between the set of VS-patterns
and the set of ROI acv-curves
was detected



This structural similarity was quantified using a functional from *partitional-clustering*



The k-partition constitutes a valid description of single-trial variability, in the post-stimulus period, for the acv-curves

1

averaged signal vs.


single-trial signals

- Bridging the gab, by means of subaveraging
- The average can be modeled as weighted average of the VQ-based prototypes

$$\overline{\mathbf{x}}(t) = \frac{1}{N} \sum_{i}^{N} \mathbf{x}_{i}(t) = \sum_{j}^{k} \alpha_{j} \mathbf{y}_{j}(t)$$

$$y_{j}(t) = \frac{\sum_{i=1}^{N} u_{ij} x_{i}(t)}{N_{j}}$$
, $\alpha_{j} = \frac{N_{j}}{N}$, $N_{j} = \sum_{i=1}^{N} u_{ij}$


$$\overline{\mathbf{x}}(t) = \frac{1}{N} \sum_{i}^{N} \mathbf{x}_{i}(t) = \sum_{j}^{k} \alpha_{j} \mathbf{y}_{j}(t)$$



This kind of modeling is informative for describing the variability in a set of responses only if the formed subaverages are different from each other



if the groups of single-trials are forming compact, well-separated clusters



Normalized scattering random partition $\rightarrow 1$




The k-partition which had been computed from the N70m-snapshots was applied to the regional and local responses.

The induced-grouping was validated, using the nSC-index, for both the set of responses.

We can consider the computed nSc-curve as the "autocorrelation" function of the N70m-variability for the Virtual-Sensor signals



as the "transfer function" showing how/if the N70m-variability manifests itself in the set of V1-ROI signals.




For the VS-signals (i.e. regional brain activity), the k-partition is a valid description well-before the stimulus-onset.

The ongoing activity before the stimulus-onset is functionally coupled with the subsequent regional response


For the V1-ROI signals (i.e. local brain activity) the k-partition is a valid description for a confined latency-range in the post-stimulus period.


There is a strong coupling of the responses in early visual areas to the regional dynamics.

Distinct response dynamics are fused into one single timecourse by ensemble-averaging




There is a loss of information; which can be restored by VQ-based grouping





Directed / Supervised Data Mining



Validation of the k-partition for the multichannel MEG signal

Concluding Remarks

The broad spectrum of N70m responses

was described by means of principal modes

which summarized the variability in the MEG signal

and further used for exploring the source of that variability

based on estimates of reconstructed brain activity

Our results revealed complex nonlinear interactions

between the sensory input and the ongoing activity

resulting to a state-dependent response

with a wide range of characteristic time scales

At the level of single-trial activity,

the polymorphic response to a simple stimulus

is generated by a coupling of polymodal areas

and cooperative activity in striate and extrastriate areas

Discussion

[Kisley & Gerstein, J. Neurosci., 1999]:

The response in rat auditory cortex is modulated by the preceding ongoing activity.

"The observed variability was attributed to rapid transitions

between different levels of thalamocortical excitability"

[Fries et al, Nat. Neurosci., 2001]:

Stimulus-induced covariation of distinct brain areas (in cats). "Coherent oscillations mediate the functional coupling between brain areas"

[Hupe et al, J.Neurophysiol., 2001]:

Feedback connections act on the early part of the responses in monkey visual cortex.

Temporally compact" visual cortex

[Lutz et al., PNAS, 2002]:

The human EEG response is modulated by background activity.

"Variability is controlled by subject's cognitive "context"

and characteristic patterns of synchrony between brain areas"



is ensemble averaging an appropriate response-recovery technique?

due to Trial-to-Trial Non-Stationarity:

Unless there is a predominant mode, the computed average is a mirage of the response generation process where different dynamics have been collapsed.






- Ordered code-book.

"The concepts I understand are great,
but I believe that the concepts I can't understand are great too.
However, the reader needs to be an excellent swimmer
like those from Dilos, so not to be drown from this (code) book ".

- Socrates

With many thanks



to a lot of people